I J C R M
International Journal of Contemporary Research In Multidisciplinary

Research Article

A Study of Set Operations and Their Applications in Mathematics

Vijay Shriram Argade
Assistant Professor, Department of Mathematics, Balbhim Arts, Science and Commerce College, Beed, Maharashtra, India

Corresponding Author: * Vijay Shriram Argade

DOI: https://doi.org/10.5281/zenodo.24273800

ABSTRACT

This article examines the principal operations performed on sets and explains how they support mathematical reasoning in several areas. A set provides a precise way to describe a collection whose membership can be determined unambiguously. Starting from this idea, the paper discusses union, intersection, set difference, complement and symmetric difference, together with their basic identities. Finite examples are used to illustrate the calculations, while Venn diagrams offer a visual interpretation of membership. The discussion then connects set operations with counting, logic, probability, relations, functions, algebra, analysis, topology, geometry and computing. The paper is an expository study based on established mathematical concepts; its purpose is to organise these ideas and demonstrate their usefulness rather than to claim a new theorem.

Manuscript Information

- **ISSN No:** 2583-7397
- **Received:** 16-09-2026
- **Accepted:** 08-11-2026
- **Published:** 16-11-2026
- **IJCRM:**5(6); 2026: 209-213
- **©2026, All Rights Reserved**
- **Plagiarism Checked:** Yes
- **Peer Review Process:** Yes

How to Cite this Article

Argade VS. A Study of Set Operations and Their Applications in Mathematics. Int J Contemp Res Multidiscip. 2026;5(6):209-213.

Access this Article Online

www.multiarticlesjournal.com

Keywords: elementary set theory, set operations, union, intersection, set difference, complement, symmetric difference, cardinality, inclusion–exclusion, mathematical logic.

© 2026 Vijay Shriram Argade. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY NC ND).https://creativecommons.org/licenses/by/4.0/

1. INTRODUCTION

Mathematical investigations frequently begin by specifying the objects under consideration. These objects may be numbers satisfying an inequality, solutions of an equation, points in a region or outcomes of an experiment. Set theory supplies a common notation for describing such collections and for expressing relationships between them. It is therefore used throughout both elementary and advanced mathematics.

A set is a collection of distinct objects for which membership is clearly determined. The objects are called elements or members. Capital letters such as A, B and C commonly denote sets. The expression $x \in A$ states that x belongs to A, whereas $x \notin A$ states that it does not. Since a set is determined by its members, the order in which they are written and repeated listings do not change the set.

When a problem involves more than one collection, set operations provide a systematic way to combine or compare them. Union describes membership in at least one set; intersection selects members shared by the sets; difference retains members of one set after excluding those in another; and complement records the members of a specified universal set that are not in the chosen set. Symmetric difference selects members that occur in exactly one of two sets.

2. Objectives and Scope of the Study

This paper has six related aims: to review the notation used for sets; to define and illustrate the main operations; to examine their standard algebraic properties; to interpret them through Venn diagrams and finite examples; to explain the role of cardinality and inclusion–exclusion; and to outline their use across mathematical subjects. The treatment is descriptive and analytical, with emphasis on classical set theory and finite collections. Connections with selected advanced topics are included to show the wider relevance of the subject.

3. Basic Concepts and Notation

3.1 Representation of Sets

A set may be described by listing its members or by stating the condition that they satisfy. The listing approach is called roster form; for instance, $A = \{2, 4, 6, 8\}$. In set-builder notation, the same collection can be written as $A = \{x \in N : x \text{ is even and } x \leq 8\}$. Roster notation is convenient when the set is small, whereas set-builder notation is often clearer for large or infinite collections.

3.2 Types of Sets

A finite set has a fixed, countable number of elements, while an infinite set has no finite number of elements. The empty set, written $\emptyset$, has no members; a singleton has precisely one. Two sets are equal when each has exactly the members of the other. Sets are disjoint if they have no common element, equivalently $A \cap B = \emptyset$. The notation $A \subseteq B$ means every member of A is also a member of B; if A is contained in B but is not equal to B, it is a proper subset.

3.3 Universal Set and Power Set

A universal set U specifies the overall domain relevant to a particular discussion. Complements are always taken relative to this chosen domain. The power set P(A) consists of every subset of A, including $\emptyset$ and A itself. For a finite set with n distinct elements, the number of subsets is 2^n. For example, when $A = \{a, b\}$, its power set is $\{\emptyset, \{a\}, \{b\}, \{a, b\}\}$, which contains four subsets.

3.4 Equality of Sets

Set equality depends on membership alone. Consequently, changing the order of the elements does not produce a different set: $\{1, 2, 3\} = \{3, 2, 1\}$. Repeating an element also has no effect, so $\{1, 1, 2, 3\}$ represents the same set as $\{1, 2, 3\}$. This is an important distinction between sets and ordered collections such as sequences.

4 Union of Sets

For sets A and B, their union contains every element that belongs to A, to B, or to both. The notation is $A \cup B = \{x : x \in A \text{ or } x \in B\}$. Here, "or" is inclusive: an element common to both sets is still included, but it is written only once.

Example: Let $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$. Then A □ B = {1, 2, 3, 4, 5, 6}. The common elements 3 and 4 are not repeated.

Union is appropriate when membership in either of two categories is sufficient. For example, if A contains students enrolled in Mathematics and B contains students enrolled in Statistics, $A \cup B$ represents those enrolled in at least one of the two subjects, including students taking both.

Some immediate union identities are $A \cup \emptyset = A$, $A \cup A = A$ and $A \cup U = U$. The operation is commutative, so $A \cup B = B \cup A$. The same idea extends to a family of sets by collecting all elements that occur in at least one member of that family.

5. Intersection of Sets

The intersection $A \cap B$ is formed from precisely those elements that are members of both A and B. In set-builder notation, $A \cap B = \{x: x \in A \text{ and } x \in B\}$. Thus, unlike union, intersection requires both membership conditions to hold.

Example: If $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$, then A □ B = {3, 4}. These are the only elements that occur in both sets.

Two sets are called disjoint when their intersection is empty. The even integers and odd integers, for example, have no common members. Intersection is useful whenever a mathematical question imposes simultaneous conditions; it selects the objects that meet every condition being considered.

The standard intersection identities include $A \cap U = A$, $A \cap \emptyset = \emptyset$, and $A \cap A = A$. Intersection is commutative and associative. Associativity means that the placement of parentheses does not affect the intersection of three or more sets.

6. Difference of Sets

The relative difference $A - B$ consists of those elements that are in A but are not in B. It can also be written A \\ B and defined by

© 2026 Vijay Shriram Argade. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY NC ND).https://creativecommons.org/licenses/by/4.0/

$A - B = \{x: x \in A \text{ and } x \notin B\}$. The order of the sets matters, so difference is not generally commutative.

Example: Let $A = \{1, 2, 3, 4, 5\}$ and $B = \{4, 5, 6, 7\}$. Then $A \setminus B = \{1, 2, 3\}$, whereas $B \setminus A = \{6, 7\}$. Hence $A \setminus B \neq B \setminus A$.

Difference can be represented through intersection with a complement: $A - B = A \cap B^c$, where B^c is taken relative to the universal set. This form is useful when simplifying expressions. In applications, it describes members of one category after excluding those that also belong to a second category.

7. Complement of a Set

Let U be the universal set and suppose $A \subseteq U$. The complement of A, denoted A^c (or A'), contains exactly those members of U that are not in A. In symbols, $A^c = \{x \in U: x \notin A\}$. The choice of U is essential because changing the universal set may change the complement.

Example: If $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ and $A = \{2, 4, 6, 8, 10\}$, then $A = \{1, 3, 5, 7, 9\}$. The complement contains the elements of U that are absent from A.

Complementation satisfies $(A^c)^c = A$, $A \cup A^c = U$ and $A \cap A^c = \emptyset$. In addition, $U^c = \emptyset$ and $\emptyset^c = U$. These identities are widely used when expressing the negation of a condition or the non-occurrence of an event.

8. Symmetric Difference

The symmetric difference of A and B contains elements belonging to one of the sets but not to both. It is denoted by $A \triangle B$ and may be defined as $(A - B) \cup (B - A)$. Equivalently, $A \triangle B = (A \cup B) - (A \cap B)$.

For $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$, the elements exclusive to A are $A - B = \{1, 2\}$, and those exclusive to Bare $B - A = \{5, 6\}$. Hence $A \triangle B = \{1, 2, 5, 6\}$; the common elements 3 and 4 are not included.

Symmetric difference is commutative and associative. It also satisfies $A \triangle \emptyset = A$ and $A \triangle A = \emptyset$. The operation is useful for comparing two collections: it isolates what is present in one collection but absent from the other. For instance, it describes students taking exactly one of two optional courses.

9. Algebraic Laws of Set Operations

The operations of union and intersection obey algebraic identities that allow set expressions to be transformed without changing their members. These laws are useful in proofs, simplification and Boolean algebra. In the identities below, A, B and C are understood to be subsets of the same universal set U.

Law	Union	Intersection
Commutative	$A \cup B = B \cup A$	$A \cap B = B \cap A$
Associative	$(A \cup B) \cup C = A \cup (B \cup C)$	$(A \cap B) \cap C = A \cap (B \cap C)$
Distributive	$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$	$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
Identity	$A \cup \emptyset = A$	$A \cap U = A$
Idempotent	$A \cup A = A$	$A \cap A = A$
Domination	$A \cup U = U$	$A \cap \emptyset = \emptyset$

9.1 De Morgan's Laws

De Morgan's identities state that $(A \cup B)^c = A^c \cap B^c$ and $(A \cap B)^c = A^c \cup B^c$. The first says that an element lies outside both sets precisely when it lies outside their union. The second says that an element fails to lie in the intersection when it is absent from at least one of the sets.

10. Venn Diagrams and Visual Interpretation

A Venn diagram represents sets by closed curves drawn within a rectangle that denotes the universal set. For two sets, the overlapping region displays their intersection, while the complete area covered by either circle represents their union. The portion of one circle outside the other depicts relative difference, and the region in the rectangle but outside a circle represents a complement.

These diagrams make membership relationships easier to interpret by turning symbolic operations into regions. They can show whether an element belongs to the first set, the second set, both or neither. A diagram with three sets has eight distinct membership regions, corresponding to the possible combinations of membership and non-membership.

10.1 Interpreting Two-Set Diagrams

For two sets, the universal region can be separated into four disjoint parts: members of A only, members of B only, members common to A and B, and members of neither set. These regions are represented respectively by $A - B$, $B - A$, $A \cap B$ and $(A \cup B)^c$.

10.2 Limits of Venn Diagrams

Venn diagrams are most convenient when only a few sets are involved. As the number of sets increases, the drawing becomes harder to construct and read. A diagram can support intuition, but it is not a substitute for a proof: a formal argument should establish that an arbitrary element belongs to one side of an identity exactly when it belongs to the other.

11. Cardinality and Counting Principles

For a finite set A, its cardinality is the number of distinct elements it contains, written $|A|$ or $n(A)$. For example, if $A = \{a, b, c, d\}$, then $|A| = 4$. Cardinality links set operations with counting and provides a way to calculate the size of unions.

For two finite sets, the inclusion–exclusion formula is $|A \cup B| = |A| + |B| - |A \cap B|$. The common elements must be subtracted once because they are included in both $|A|$ and $|B|$ when the two cardinalities are added.

For three finite sets, the corresponding formula is $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$. Pairwise intersections are first subtracted to correct double counting; the triple intersection is then added back because it has otherwise been removed too many times.

Suppose 28 students study Mathematics, 20 study Statistics and 9 study both subjects. The number studying at least one subject is $28 + 20 - 9 = 39$. This use of inclusion–exclusion assumes that all three figures refer to the same group of students.

© 2026 Vijay Shriram Argade. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY NC ND).https://creativecommons.org/licenses/by/4.0/

12. Worked Mathematical Examples

Example 1: Union and intersection

Let A = {2, 4, 6, 8} and B = {4, 8, 10, 12}. The union contains every element appearing in either set: A □ B = {2, 4, 6, 8, 10, 12}. The intersection contains only the common elements: A □ B = {4, 8}.

Example 2: Difference

Let P = {a, b, c, d, e} and Q = {c, d, f}. Then P □ Q = {a, b, e}, while Q □ P = {f}. The result demonstrates that set difference is not commutative.

Example 3: Complement

Take U = {1, 2, 3, 4, 5, 6, 7, 8} and M = {1, 3, 5, 7}. Relative to U, the complement is M^c = {2, 4, 6, 8}. It follows that M ∪ M^c = U and M ∩ M^c = ∅.

Example 4: Cardinality

Suppose |A| = 32, |B| = 24 and |A □ B| = 10. Then |A □ B| = 32 + 24 □ 10 = 46. Therefore, 46 elements belong to at least one of the sets.

Example 5: Symmetric difference

Let X = {1, 2, 3, 4} and Y = {3, 4, 5, 6}. The elements exclusive to X are {1, 2}, and those exclusive to Y are {5, 6}.

Thus, X Y = {1, 2, 5, 6}.

Example 6: Applying De Morgan's law

Let U = {1, 2, 3, 4, 5, 6}, A = {1, 2, 3} and B = {3, 4}. Then A ∪ B = {1, 2, 3, 4}, so $(A \cup B)^c$ = {5, 6}. Also, A^c = {4, 5, 6} and B^c = {1, 2, 5, 6}; therefore $A^c \cap B^c$ = {5, 6}. The two sides agree, verifying this instance of De Morgan's first law.

13. Applications in Mathematical Logic

Set operations have direct logical counterparts. Union corresponds to inclusive disjunction (OR), intersection to conjunction (AND), and complement to negation (NOT). If A is the collection of objects for which proposition P is true and B is the collection for which Q is true, then A ∩ B describes objects satisfying both propositions, while A ∪ B describes objects satisfying at least one.

The set identities of De Morgan correspond to familiar logical equivalences: the negation of P OR Q is (not P) AND (not Q), while the negation of P AND Q is (not P) OR (not Q). This connection is useful in predicate logic, mathematical proofs and the simplification of compound statements.

14. Applications in Probability Theory

In probability, the sample space S is the set of all possible outcomes, and an event is a subset of S. For events A and B, the union A ∪ B occurs when at least one of the events occurs; the intersection A ∩ B occurs when both occur. The complement A^c represents the event that A does not occur.

The addition rule for two events is P(A ∪ B) = P(A) + P(B) − P(A ∩ B), which mirrors the inclusion–exclusion formula for finite sets. If the events are mutually exclusive, their intersection is empty and the probability of their union is the sum of their probabilities. The complement rule is $P(A^c) = 1 - P(A)$.

Consider one throw of a fair six-sided die. Let A = {2, 4, 6} be the event of obtaining an even result and B = {4, 5, 6} the event of obtaining a result greater than 3. Then A ∩ B = {4, 6}, whereas A ∪ B = {2, 4, 5, 6}. This example shows how intersections and unions describe combined events.

15. Applications in Relations and Functions

A relation from A to B is a subset of the Cartesian product A × B, whose elements are ordered pairs (a, b) with a ∈ A and b ∈ B. Since relations are sets of ordered pairs, set operations can be used to compare or combine relations. In particular, the union of two relations from A to B is also a relation from A to B.

A function is a relation with the additional requirement that each domain element is assigned exactly one output. Its domain and codomain are sets, and its range is contained in the codomain. Inverse images are also sets: under a function, inverse images preserve unions, intersections and complements when complements are taken relative to the corresponding domain and codomain.

16. Applications in Algebra and Number Theory

Number systems are naturally expressed as nested sets. With the convention that natural numbers are included in the integers, the usual containment sequence is N ⊆ Z ⊆ Q ⊆ R ⊆ C, where these symbols denote the natural, integer, rational, real and complex numbers, respectively. Each inclusion indicates that every member of one system also belongs to the next.

In algebra, a solution set records all values satisfying a specified condition. If A and B are solution sets for two conditions, A ∩ B contains the values satisfying both conditions, while A ∪ B contains those satisfying at least one. For example, the simultaneous inequalities x > 1 and x < 5 have solution set $(1, \infty) \cap (-\infty, 5) = (1, 5)$.

17. Applications in Real Analysis and Topology

Real analysis uses sets to describe domains, ranges, intervals and neighbourhoods associated with functions and sequences. For example, the open interval (a, b) is the set of real numbers x for which a < x < b. Unions and intersections of intervals provide a concise way to specify more complicated subsets of the real line.

Topology studies a space through a selected family of subsets called open sets. One defining property is that an arbitrary union of open sets is open and a finite intersection of open sets is open. For closed sets, arbitrary intersections and finite unions remain closed. These properties demonstrate the central role of set operations in topology.

18. Applications in Geometry

Geometric objects can be treated as sets of points in the plane or in three-dimensional space. Under this interpretation, the intersection of two figures gives their common points, their union gives all points belonging to either figure, and their difference selects points in one figure but not the other.

For example, let A be the set of points in a square and B the set of points in a circular region. The set A ∩ B consists of points common to the two regions. Such descriptions are useful in coordinate geometry, geometric modelling and the analysis of plane or spatial regions.

© 2026 Vijay Shriram Argade. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY NC ND).https://creativecommons.org/licenses/by/4.0/

19. Applications in Discrete Mathematics and Computing

Sets are basic objects in discrete mathematics, including combinatorics, graph theory and algorithm design. A graph, for example, is specified through a set of vertices and a set of edges. Comparing vertex sets can reveal shared or distinct vertices. In programming, set-based data structures support membership tests and operations such as union, intersection and difference.

Database queries also involve operations resembling those of set theory. Results from separate queries may be combined or compared to identify common records or records appearing in only one result. Care is needed, however, because many database systems retain duplicate rows by default; such behaviour is closer to a multiset model than to the classical mathematical definition of a set.

20. RESEARCH METHODOLOGY

The paper uses a descriptive and analytical approach. It first presents standard definitions and notation for sets and their operations. It then organises key algebraic identities, illustrates them through finite examples and interprets selected results using Venn diagrams. The final part relates the concepts to several branches of mathematics and computing.

The discussion is based on established mathematical texts and does not claim a new theorem, experiment or original dataset. Its contribution is a consolidated educational account that places definitions, identities, worked calculations and applications within one framework. The examples are deliberately small so that each result can be checked directly.

21. RESULTS AND DISCUSSION

The comparison of the five operations highlights their different functions: union collects elements from either set, intersection selects shared members, difference removes the members of a second set from the first, complement selects elements outside a set within a stated universe, and symmetric difference retains elements belonging to exactly one of the two sets.

The algebraic identities provide a consistent framework for manipulating set expressions. Commutativity and associativity permit changes in order and grouping; distributivity connects union with intersection; and complement identities, including De Morgan's laws, describe how operations behave under negation. Inclusion–exclusion extends these ideas to finite counting.

The applications reviewed here show that set operations are used well beyond introductory set theory. They describe logical conditions and probability events, support the formal definition of relations and functions, express the containment of number systems, and help formulate concepts in analysis, topology, geometry and discrete mathematics.

Several points deserve particular attention when applying these operations. Relative difference depends on the order of the sets; a complement is undefined until the universal set is specified; a union lists a common element only once; and the word "or" in the definition of union includes the case in which both conditions hold. These distinctions prevent frequent errors in calculations.

22. CONCLUSION

Set operations offer a concise and rigorous language for comparing and combining mathematical collections. The five operations considered in this paper capture different forms of membership, while their algebraic laws make it possible to simplify expressions and establish identities. Venn diagrams provide an accessible visual aid, and cardinality together with inclusion–exclusion connects the subject to finite counting and probability. The examples from logic, functions, algebra, analysis, topology, geometry and computing demonstrate the broad reach of these ideas. A sound command of set operations therefore supports further study across many branches of mathematics.

The standard definitions and properties discussed in this paper are treated in classical texts on set theory, discrete mathematics and topology (Halmos, 1960; Enderton, 1977; Rosen, 2019; Munkres, 2000).

REFERENCES

1. Halmos PR. *Naive set theory*. Princeton (NJ): D. Van Nostrand Company; 1960.
2. Enderton HB. *Elements of set theory*. New York: Academic Press; 1977.
3. Rosen KH. *Discrete mathematics and its applications*. 8th ed. New York: McGraw-Hill Education; 2019.
4. Lipschutz S, Lipson M. *Schaum's outline of set theory and related topics*. 2nd ed. New York: McGraw-Hill; 2009.
5. Devlin K. *The joy of sets: fundamentals of contemporary set theory*. 2nd ed. New York: Springer; 1993.
6. Munkres JR. *Topology*. 2nd ed. Upper Saddle River (NJ): Prentice Hall; 2000.

Creative Commons (CC) License

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution–Non-Commercial–No Derivatives 4.0 International (CC BY-NC-ND 4.0) license. This license permits sharing and redistribution of the article in any medium or format for non-commercial purposes only, provided that appropriate credit is given to the original author(s) and source. No modifications, adaptations, or derivative works are permitted under this license.

About the Author



Vijay Shriram Argade is an Assistant Professor in the Department of Mathematics at Balbhim Arts, Science and Commerce College, Beed, Maharashtra, India. His academic interests include mathematics, mathematical analysis, and contemporary developments in mathematical research. He is dedicated to teaching, research, and fostering analytical and problem-solving skills among students.

© 2026 Vijay Shriram Argade. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY NC ND).https://creativecommons.org/licenses/by/4.0/